

Electron Scattering off ^4He with Three-Nucleon Forces

Sonia Bacca

TRIUMF, 4004 Wesbrook Mall, Vancouver, B.C. V6J 2A3, Canada

Nir Barnea

Racah Institute of Physics, Hebrew University, 91904, Jerusalem, Israel

Winfried Leidemann, Giuseppina Orlandini

*Dipartimento di Fisica, Università di Trento and INFN (Gruppo Collegato di Trento),
via Sommarive 14, I-38100 Trento, Italy*

Abstract

An *ab initio* calculation of the ^4He (e,e') inelastic longitudinal response function R_L is presented. Realistic two- and three-body forces are used. The four-body continuum dynamics is treated rigorously with the help of the Lorentz integral transform. The three-nucleon force reduces the quasi-elastic peak height by about 10% for momentum transfers q between 300 and 500 MeV/c. Experimental data are well described, but not sufficiently precise to resolve this effect. The reduction due to the three-nucleon force increases significantly at lower q reaching up to about 40% at $q = 100$ MeV/c. However, at such q values data are missing.

Key words: three-nucleon force, electron scattering, Lorentz integral transform

PACS: 21.45-v, 21.45.Ff, 25.30Fj

The longitudinal response function R_L is given by

$$R_L(\omega, q) = \sum_f |\langle \Psi_f | \hat{\rho}(q) | \Psi_0 \rangle|^2 \delta \left(E_f + \frac{q^2}{2M} - E_0 - \omega \right), \quad (1)$$

where M is the target mass, $|\Psi_{0/f}\rangle$ and $E_{0/f}$ denote the four-body initial and final state wave functions and energies, respectively, while ω and q are the energy and momentum transfers. The charge density operator ρ is defined as

$$\hat{\rho}(q) = \frac{e}{2} \sum_i (1 + \tau_i^3) \exp[i\mathbf{q} \cdot \mathbf{r}_i], \quad (2)$$




Fig. 1. $R_L(\omega, q)$ at various q . Dashed lines: AV18; solid lines: AV18+UIX. Data from Bates [9] (squares), Saclay [10] (circles) and world data-set from [11] (triangles).

where e is the proton charge and τ_i^3 the isospin third component of nucleon i . For a conventional calculation of R_L one would need to know explicitly the four-body continuum state wave functions Ψ_f . In the Lorentz integral transform (LIT) method [1] this difficulty is circumvented by considering instead of $R_L(\omega, q)$ an integral transform $\mathcal{L}_L(\sigma, q)$ with a Lorentzian kernel defined for a complex parameter $\sigma = \sigma_R + i\sigma_I$, which is then inverted in order to obtain $R_L(\omega, q)$ (see review [2]).

For the calculation of R_L we take a realistic nuclear interaction consisting in the AV18 two-nucleon potential [3] and the UIX three-nucleon force (3NF) [4]. The ^4He ground-state wave function and the LIT are calculated using expansions in hyperspherical harmonics with the EIH [5,6] and Lanczos [7] techniques (for more information concerning the calculation see [8]).

In Fig. 1 we show R_L at $300 \text{ MeV}/c \leq q \leq 500 \text{ MeV}/c$. One sees that the 3NF reduces the quasi-elastic peak strength by about 10%. Experimental data are described quite well by our full result, but they are not precise enough to resolve the 3NF effect. In Fig. 2 we illustrate R_L at lower q . One readily notes a very strong reduction at lower energies due to the 3NF, which reaches up to about 40%. The reduction cannot be attributed to a simple binding effect as becomes evident from the also shown R_L results with a semirealistic NN force (MT potential [12]). In fact, ^4He binding energies are 24.3, 28.4, and 30.6 MeV for AV18, AV18+UIX, and MT potentials, respectively. Even though the MT energy is closer to that of AV18+UIX, the MT R_L is more similar to the R_L of AV18




Fig. 2. $R_L(\omega, q)$ at various q : AV18 (dashed), AV18+UIX (solid), MT (dash-dotted). Data in (a) from [13].

than to the AV18+UIX R_L . At lower q there is only one data set at 200 MeV/c [13], which is not sufficiently precise to draw concrete conclusions.

The work of S.B. was supported by the Natural Sciences and Engineering Research Council (NSERC) and by the National Research Council of Canada and the work of N.B. by the Israel Science Foundation (grant no. 361/05). Numerical calculations were performed at CINECA (Bologna).

References

- [1] V. D. Efros, W. Leidemann, and G. Orlandini, Phys. Lett. B **338**, 130 (1994).
- [2] V. D. Efros, W. Leidemann, G. Orlandini, and N. Barnea, J. Phys. G: Nucl. Part. Phys. **34**, R459 (2007).
- [3] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, Phys. Rev. C **51**, 38 (1995).
- [4] B. S. Pudliner, V. R. Pandharipande, J. Carlson, S. C. Pieper, and R. B. Wiringa, Phys. Rev. C **56**, 1720 (1997).
- [5] N. Barnea, W. Leidemann, and G. Orlandini, Phys. Rev. C **61**, 054001 (2000); Nucl. Phys. **A693**, 565 (2001).
- [6] N. Barnea, V. D. Efros, W. Leidemann, and G. Orlandini, Few-Body Syst. **35**, 155 (2004).
- [7] M. A. Marchisio, N. Barnea, W. Leidemann, and G. Orlandini, Few-Body Syst. **33**, 259 (2003).
- [8] S. Bacca, N. Barnea, W. Leidemann, and G. Orlandini, arXiv:0811.4624 [nucl-th].
- [9] S. A. Dytman *et al.*, Phys. Rev. C **38**, 800 (1988).
- [10] A. Zghiche *et al.*, Nucl. Phys. **A572**, 513 (1994).
- [11] J. Carlson, J. Jourdan, R. Schiavilla, and I. Sick, Phys. Rev. C **65**, 024002 (2002).
- [12] B. S. Pudliner, V. R. Pandharipande, J. Carlson, S. C. Pieper, and R. B. Wiringa, Phys. Rev. C **56**, 1720 (1997).
- [13] A. Yu. Buki, I. S. Timchenko, N. G. Shevchemko, and I. A. Nenko, Phys. Lett. B **641**, 156 (2006).